


WATER AND SANITATION

Benefits and costs of drinking water and sanitation interventions in Andhra Pradesh



Cost-Benefit Analysis

AUTHORS: Bjorn
Larsen
Environmental Economist,
Independent Consultant

© 2018 Copenhagen Consensus Center

info@copenhagenconsensus.com

www.copenhagenconsensus.com

This work has been produced as a part of the Andhra Pradesh Priorities project under the larger, India Consensus project.

This project is undertaken in partnership with Tata Trusts.

TATA TRUSTS

Some rights reserved



This work is available under the Creative Commons Attribution 4.0 International license ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)). Under the Creative Commons Attribution license, you are free to copy, distribute, transmit, and adapt this work, including for commercial purposes, under the following conditions:

Attribution

Please cite the work as follows: #AUTHOR NAME#, #PAPER TITLE#, Andhra Pradesh Priorities, Copenhagen Consensus Center, 2017. License: Creative Commons Attribution CC BY 4.0.

Third-party content

Copenhagen Consensus Center does not necessarily own each component of the content contained within the work. If you wish to re-use a component of the work, it is your responsibility to determine whether permission is needed for that re-use and to obtain permission from the copyright owner. Examples of components can include, but are not limited to, tables, figures, or images.

Benefits and Costs of Drinking Water and Sanitation Interventions in Andhra Pradesh

Andhra Pradesh Priorities An India Consensus Prioritization Project

Bjorn Larsen

Environmental Economist

Independent Consultant

Working paper as of May 25, 2018

ACADEMIC ABSTRACT	1
POLICY ABSTRACT	2
THE PROBLEM	2
INTERVENTION 1: IMPROVED DRINKING WATER SOURCE	3
<i>Overview</i>	3
<i>Implementation Considerations</i>	4
<i>Costs and Benefits</i>	4
Costs.....	4
Benefits	4
INTERVENTION 2: HOUSEHOLD POINT-OF-USE TREATMENT OF DRINKING WATER	5
<i>Overview</i>	5
<i>Implementation Considerations</i>	5
<i>Costs and Benefits</i>	5
Costs.....	5
Benefits	5
INTERVENTION 3: IMPROVED SANITATION	6
<i>Overview</i>	6
<i>Implementation Considerations</i>	6
<i>Costs and Benefits</i>	6
Costs.....	6
Benefits	6
INTERVENTION 4: PROMOTION OF THE USE OF SANITATION FACILITIES	7
<i>Overview</i>	7
<i>Implementation Considerations</i>	7
<i>Costs and Benefits</i>	7
Costs.....	7
Benefits	8
BCR TABLE	8
1. INTRODUCTION	10
1.1 CONTEXT AND INTERVENTIONS	10
1.2 COMMON DATA	10
1.3 LITERATURE REVIEW	12
1.4 HOUSEHOLD DRINKING WATER AND SANITATION SITUATION	14
2. IMPROVED DRINKING WATER SOURCE	15
2.1 DESCRIPTION OF INTERVENTION.....	15
2.2 CALCULATION OF COSTS AND BENEFITS.....	16
2.2.1 <i>Costs</i>	16

2.2.2 Benefits.....	16
2.2.3 Benefit-cost ratios.....	19
2.3 ASSESSMENT OF QUALITY OF EVIDENCE	20
2.4 SENSITIVITY ANALYSIS	20
3. HOUSEHOLD POINT-OF-USE TREATMENT OF DRINKING WATER.....	21
3.1 DESCRIPTION OF INTERVENTION.....	21
3.2 LITERATURE REVIEW	22
3.3 CALCULATION OF COSTS AND BENEFITS.....	22
3.3.1 Costs.....	23
3.3.2 Benefits.....	23
3.3.3 Benefit-cost ratios.....	25
3.4 ASSESSMENT OF QUALITY OF EVIDENCE	26
3.5 SENSITIVITY ANALYSIS	27
4. IMPROVED SANITATION.....	28
4.1 DESCRIPTION OF INTERVENTION.....	28
4.2 CALCULATION OF COSTS AND BENEFITS.....	29
4.2.1 Costs.....	29
4.2.2 Benefits.....	30
4.2.3 Benefit-cost ratios.....	33
4.3 ASSESSMENT OF QUALITY OF EVIDENCE	34
4.4 SENSITIVITY ANALYSIS	35
5. PROMOTION OF THE USE OF SANITATION FACILITIES	36
5.1 DESCRIPTION OF INTERVENTION.....	36
5.2 LITERATURE REVIEW	36
5.3 CALCULATION OF COSTS AND BENEFITS.....	36
5.3.1 Costs.....	36
5.3.2 Benefits.....	38
5.3.3 Benefit-cost ratios.....	40
5.4 ASSESSMENT OF QUALITY OF EVIDENCE	41
6. CONCLUSION.....	42
REFERENCES	44
ANNEX 1. HEALTH EFFECTS OF UNSAFE DRINKING WATER AND SANITATION.....	46

Academic Abstract

This paper evaluates the benefits and costs of four household drinking water and sanitation interventions. The interventions were selected on the basis of coverage rates and behavioral dimensions in the state. Benefits and costs are presented as a ratio of annualized benefits and annualized costs (benefit-cost ratios) over the expected useful life of each intervention.

Benefit-cost ratios (BCRs) are found to be the largest for improved rural sanitation (7.0-8.3) and improved urban water supply (6.8-10.7), followed by improved urban sanitation (5.7-6.4) household point-of-use (POU) treatment of drinking water (6.5-6.9) and improved rural water supply (3.2-5.0). A behavioral change campaign (BCC) promoting the use of existing sanitation by all household members has the lowest BCR (1.7-1.9).

These BCRs reflect benefits estimated using a value of statistical life (VSL) for averted deaths, a monetary value of time savings at 50-100% of wage rates, the cost-of-illness (COI) approach for averted illness, and annual discount rates of 3-8%. BCRs with averted years of life lost to mortality (YLL) and illness (YLD), valued at 3 times GDP per capita, are also presented in the report. The latter BCRs are somewhat lower than the former, especially at higher discount rates.

The quality of evidence associated with the estimated benefits and costs of these interventions range from “medium” to “medium-strong”. The BCC campaign is, however, supported by only “limited” quality of evidence.

Policy Abstract

The Problem

Diarrheal disease caused nearly 1.7 million deaths globally in 2016. Nearly 0.8 million of these deaths occurred in India of which 90% were due to unsafe drinking water, sanitation and hygiene according to estimates by the Global Burden of Disease 2016 (GBD 2016).¹

In Andhra Pradesh over 30 thousand deaths were caused by diarrheal disease and intestinal infections (mainly typhoid and paratyphoid) in 2016, constituting 8.0% of all deaths in the state according to GBD 2016, thus remaining a major cause of concern in relation to drinking water and sanitation.

About 73% of households in Andhra Pradesh had access to an improved drinking water source in 2015-16 according to the National Family Health Survey 4 (NFHS 4).² Improved drinking water sources are generally considered having a lesser risk of pathogen contamination than unimproved sources (e.g., shallow dug wells, unprotected springs, open surface water). Water from improved sources are, however, not necessarily safe to drink (Pruss-Ustun et al, 2014). Nevertheless, only 19% of households practiced point-of-use (POU) treatment of their drinking water in 2005-06, mainly by filtering (11%) and boiling (8%) (NFHS 3).³ Moreover, 14% of the population had more than 30 minutes round-trip to their drinking water source in 2005-06 (NFHS 3), particularly affecting women and children who often perform the task of fetching drinking water.⁴ It should be noted that these figures from the NFHS 3 pertains to Andhra Pradesh before the separation of Telengana.

About 61% of households in Andhra Pradesh had a sanitary toilet facility in 2015 according to the Swachhta Status Report 2016 (MSPI/GOI, 2017). Moreover, 54% of households had access to an improved non-shared sanitation facility in 2015-16 according to the National

¹ <http://www.healthdata.org/>

² The Joint Monitoring Programme for Water Supply and Sanitation (JMP) of WHO/UNICEF classifies household drinking water sources and sanitation facilities into improved and unimproved sources/facilities.

³ The figures for 2015-16 from the NFHS 4 were not yet publicly available at the time of this paper.

⁴ The figures for 2015-16 from the NFHS 4 were not yet publicly available at the time of this paper.

Family Health Survey IV.⁵ Thus about 7% of households shared a sanitation facility with other households in 2015.

The Swachhta Status Report 2016 also reveals that 2.2% of persons with a sanitary toilet facility continued to practice open defecation (OD). Additionally, 39% of households had no sanitation facility and practiced OD in 2015 (MSPI/GOV, 2017). A substantial drive by the government for household sanitation and eradication of OD has, however, reduced this figure quite substantially since 2015.

Based on the household drinking water and sanitation situation in the state, four interventions are evaluated in this paper in terms of their benefits and costs:

- 1) Improved drinking water source within 30 minutes round-trip from household dwelling.
- 2) Household POU treatment of drinking water with a filtering device.
- 3) Non-shared, improved sanitation facility.
- 4) Household members full use of existing sanitation facilities.

Benefits and costs are presented as a ratio of annualized benefits and annualized costs (benefit-cost ratios (BCRs)) over the expected useful life of each intervention.

It should be noted that a comparison of benefits and costs of these four interventions does not imply that the interventions are mutually exclusive. Indeed they can all contribute towards reducing the disease burden from inadequate drinking water and sanitation. However, a ranking of the interventions in terms of high to low benefit-cost ratios (BCRs) provide valuable information as to setting priorities when facing limited resources.

Intervention 1: Improved drinking water source

Overview

The most common type of improved drinking water source is piped water to dwelling in urban areas and public tap/standpipe or tubewell/borehole in rural areas (NSS 71, 2014). Thus the intervention is:

⁵ The Joint Monitoring Programme (JMP) of WHO/UNICEF classifies household sanitation into improved and unimproved facilities.

- i) Piped water to dwelling for urban households without an improved drinking water source.
- ii) Tubewell/borehole for rural households without an improved drinking water source or that have more than 30 minutes round-trip to their drinking water source.

Implementation Considerations

An implementation consideration in rural areas is the location of an improved drinking water source relative to household dwelling. A household may continue to use an unimproved source if this source is closer to the dwelling than the improved source. Providing an improved drinking water source within 30 minutes round-trip may also have cost implications if households are sparsely distributed.

Costs and Benefits

Costs

Annualized cost of improved drinking water supply is estimated in the range of Rs. 1,233 – 1,753 per household. This includes initial capital cost of piped water supply or tubewell/borehole and annual operations and maintenance cost (O&M). About 4.7 million households are expected to benefit from this intervention with total annualized cost of Rs. 5.8-8.2 billion.

Benefits

The benefits of the intervention quantified are the value of health improvements and productivity benefits. The largest benefits are avoided mortality and productivity benefits in terms of time savings from water source closer to dwelling. Total annual benefits are Rs. 21-40 billion, depending on valuation method applied. The value of averted deaths accounts for 44-64% of total benefits. Health care cost savings account for about 3-5% of total benefits. Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for about 10-15% of total benefits. And productivity benefits in terms of time savings account for 23-36% of total benefits.

Intervention 2: Household point-of-use treatment of drinking water

Overview

Only 19% of households in Andhra Pradesh (and Telangana) practiced appropriate methods of point-of-use (POU) treatment of drinking water a decade ago, compared to over 20% nationwide in India according to the NFHS 3 (2005-06). POU treatment by appropriate methods is likely to have increased somewhat, postulated at 23% currently. Given that water filter was the most common method in Andhra Pradesh, the intervention assessed in this paper is a behavioral change campaign (BCC) that promotes household POU treatment of drinking water with water filter.

Implementation Considerations

A challenge with a BCC program is decisions as to its intensity, which affects both cost and the rate of behavioral change. Thus emphasis ought to be placed on identifying and testing a design that gives a desired behavioral response at an acceptable cost.

Costs and Benefits

Costs

The main costs of POU treatment of drinking water is the water filter (Rs. 2,000), parts replacement (Rs. 500 per year), and cost of time spent on filtering water (Rs. 434 per year). For scenarios that make 1.4-2.8 million households start POU treatment by filtering of drinking water, the total annualized cost is Rs. 2-4.1 billion. Additionally, a BCC program is estimated to cost in the range of Rs. 0.1-1.7 billion annualized, depending on intensity or level of effort, prompting 1.4-2.8 million households to start POU treatment.

Benefits

The total value of the health and productivity benefits of the intervention is estimated at Rs. 7 – 31 billion, depending on valuation method applied and household response to the promotion program. Health care cost savings account for 6-13% of total benefits. Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for 10-23% of total benefits. The value of averted deaths accounts for the remaining benefits.

Intervention 3: Improved sanitation

Overview

The government has been/is undertaking a substantial drive for household sanitation and eradication of open defecation (OD) with an incentive up to Rs 12,000 to eligible households. Most households opt for a flush/pour-flush system with a single- or twin-pit. This is therefore the intervention assessed in this paper. A target of 95% household coverage with improved, non-shared sanitation is applied. The government may, however, aim for 100% coverage. The difference has no/negligible effect on the benefit-cost ratios (BCRs), unless the cost of reaching the last 5% of households escalates.

Implementation Considerations

A consideration is the 2.2% of people that continue to practice OD even after construction of a sanitation facility, according to the Swachhta Status Report 2016 (MSPI/GOI, 2017). A sanitation drive may therefore be accompanied by a sanitation use campaign.

Costs and Benefits

Costs

Costs of household sanitation include initial capital cost, periodic emptying of pit or septic tank, annual operations and maintenance (O&M) cost, cost or time for cleaning of sanitation facility, and cost of sanitation promotion programs by the government. An initial capital cost of Rs. 20,000 is applied. This is for the toilet, installation, and construction of superstructure (base, building). Emptying the pit every five years costs around Rs. 1,500. Annual O&M is assumed to be 5% of initial capital cost. Cleaning of sanitation facility is assumed to be 10 minutes per day. Sanitation program cost of Rs. 600 per household (5% of sanitation budget) that currently do not have improved, non-shared sanitation is assumed to cover promotion of household sanitation, proper designs, and promotion of household members' full use of the facility.

Annualized average rural and urban cost per household is estimated at roughly Rs. 5,600 – 8,200 and total annualized cost of intervention is estimated at Rs. 31-34 billion based on total intervention beneficiaries of 5.4 million households.

Benefits

The total value of the health and productivity benefits of the intervention is estimated at Rs. 143 – 251 billion, depending on valuation method applied. Health care cost savings account for 2-4% of total benefits. Productivity benefits in terms of time savings account for 40-61% of total benefits. Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for 6-9% of total benefits. The value of averted deaths accounts for 27-52%.

Intervention 4: Promotion of the use of sanitation facilities

Overview

The Swachhta Status Report 2016 found that 2.2% of household members continue to practice OD after construction of sanitation facility. Faced with a situation of OD among households with sanitation facility, the intervention is a behavioral change campaign (BCC) that promotes the consistent use of existing sanitation facilities.

Implementation Considerations

A challenge with a BCC program is decisions as to its intensity, which affects both cost and the rate of behavioral change. Thus emphasis ought to be placed on identifying and testing a design that gives a desired behavioral response at an acceptable cost.

Costs and Benefits

Costs

Annualized BCC program cost per household ranges from Rs. 11-12 for the “low” intensity program to Rs. 176-190 for the “high” intensity program. The program is directed at all households that currently have a sanitary toilet facility as reported by the Swachhta Status Report 2016 (MSPI/GOI, 2017). Thus total annualized program cost is in the range of Rs. 96-1,655 million.

As previously stated, some household members do not use their sanitation facility and continue to practice OD. The NSS 69 (2012) finds several reasons for this: i) no toilet superstructure; ii) lack of cleanliness/insufficient water; iii) malfunctioning of the latrine; iv) personal preferences; and v) other reasons. The two most important reasons in Andhra Pradesh were personal preferences (17%) and lack of cleanliness/insufficient water (62%).

Personal preferences for OD imply that these household members perceive some benefits of continuing OD, or cost of giving up their old practice of OD. The magnitude of this cost is unknown. It is therefore in this paper assumed to be (at least) as large as the value of time spent on OD. Moreover, that lack of cleanliness/insufficient water was reported as the second main reason for not using the toilet facility prompts the inclusion of cost of cleaning of toilet facility (or value of time spent on cleaning).

Total cost of intervention is therefore estimated at roughly Rs. 260 – 1,960 million, depending on intensity or level of effort of the promotion program and household member response rates. The cost includes program cost, cost associated with personal preferences, and cost or time used on cleaning toilet facilities.

Benefits

The total value of the health and productivity benefits of the intervention is estimated at Rs. 0.6 – 1.7 billion. Health care cost savings account for less than 2% of total benefits. Productivity benefits in terms of time savings account for 64-77% of total benefits. Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for about 4-5% of total benefits. The value of averted deaths accounts for 16-31% of total benefits.

BCR Table

The benefit-cost ratios (BCRs) are found to be the largest for improved urban drinking water supply and improved rural sanitation, followed by improved urban sanitation, household point-of-use (POU) treatment of drinking water and improved rural water supply. A behavioral change campaign (BCC) promoting the use of existing sanitation by all household members has the lowest BCR.

The quality of evidence associated with three of these interventions range from “medium” to “medium-strong”. The BCC intervention is supported by only “limited” quality of evidence.

Table 1a. Summary of the benefits and costs and interventions (Rs million annualized), VSL approach for mortality benefits and Cost of Illness approach for valuing morbidity benefits

	Interventions	Benefit	Cost	BCR	Quality of Evidence
1a	Improved drinking water supply - urban	20,294	2,280	8.9	Medium
1b	Improved drinking water supply - rural	18,167	4,403	4.1	Medium
2	BCC promotion of household POU treatment of drinking water*	23,136	3,422	6.8	Medium-Strong
3a	Improved sanitation - urban	33,527	5,441	6.2	Medium-Strong
3b	Improved sanitation - rural	207,151	26,683	7.8	Medium-Strong
4	BCC promotion of use of existing sanitation facilities*	1,192	673	1.8	Limited

Notes: All figures assume a 5% discount rate, and use VSL for valuation of mortality benefits and cost-of-illness (COI) for valuation of morbidity benefits. * “Mid” scenario. Source: Author.

Table 1b. Summary of the benefits and costs and interventions (Rs million annualized), YLL approach to valuing mortality benefits; YLD approach to measuring morbidity benefits

	Interventions	Benefit	Cost	BCR	Quality of Evidence
1a	Improved drinking water supply - urban	10,790	2,280	4.7	Medium
1b	Improved drinking water supply - rural	11,350	4,403	2.6	Medium
2	BCC promotion of household POU treatment of drinking water*	10,356	3,422	3.0	Medium-Strong
3a	Improved sanitation - urban	21,996	5,441	4.0	Medium-Strong
3b	Improved sanitation - rural	142,011	26,683	5.3	Medium-Strong
4	BCC promotion of use of existing sanitation facilities*	959	673	1.4	Limited

Notes: All figures assume a 5% discount rate, and use an approach that values discounted Years of Life Lost (YLLs) at 3x SGDP per capita for valuation of mortality benefits and values Years Lost to Disability (YLDs) also at 3x GDP per SGDP of morbidity benefits. * “Mid” scenario. Source: Author.

1. Introduction

1.1 Context and interventions

Diarrheal disease caused nearly 1.7 million deaths globally in 2016. Nearly 0.8 million of these deaths occurred in India of which 90% were due to unsafe drinking water, sanitation and hygiene according to estimates by the Global Burden of Disease 2016 (GBD 2016).

In Andhra Pradesh over 30 thousand deaths were caused by diarrheal disease and intestinal infections (mainly typhoid and paratyphoid) in 2016, constituting 8.0% of all deaths in the state according to GBD 2016, thus remaining a major cause of concern in relation to drinking water and sanitation.

Based on the household drinking water and sanitation situation in the state, four interventions are evaluated in this paper in terms of their benefits and costs:

- 1) Improved drinking water source within 30 minutes round-trip from household dwelling.
- 2) Household POU treatment of drinking water with a filtering device.
- 3) Non-shared, improved sanitation facility.
- 4) Household members full use of existing sanitation facilities.

Benefits and costs are presented as a ratio of annualized benefits and annualized costs (benefit-cost ratios (BCRs)) over the expected useful life of each intervention. BCRs are presented using discount rates of 3%, 5% and 8%.

It should be noted that a comparison of benefits and costs of these four interventions does not imply that the interventions are mutually exclusive. Indeed they can all contribute towards reducing the disease burden from inadequate drinking water and sanitation. However, a ranking of the interventions in terms of high to low benefit-cost ratios (BCRs) provide valuable information as to setting priorities when facing limited resources.

1.2 Common data

Many of the data utilized in this paper are common to the four interventions. These data are discussed in this section.

Data on household access to drinking water and sanitation, and household point-of-use (POU) treatment of drinking water are mainly from the National Family Health Surveys (NFHS) 4 (2015-16) and 3 (2005-06), and the National Sample Surveys (NSS) 71 (2014) and 69 (2012). The NFHS 4 provides state level data for Andhra Pradesh while the preceding surveys include Telengana. State level data on the use of existing sanitation facilities are from the Swachhta Status Report 2016 (MSPI/GOI, 2017). This report also provides state level data on household sanitation coverage and open defecation (OD) rates in 2015.

Assessed benefits of the interventions are health improvements and productivity benefits. Health improvements are averted cases of diarrheal and intestinal infectious disease (typhoid and paratyphoid) and associated mortality. Included are also averted cases of infectious disease mortality among young children from a reduction in the effect of repeated diarrheal infections on child underweight and associated mortality. The methodology for estimating health improvements of the interventions are provided in Annex 1.

Other health benefits, such as reduction in intestinal nematode infections, are not included in the assessment, as these health effects are relatively minor in terms of total disease burden compared to mortality and morbidity from diarrheal disease and intestinal infections (typhoid, paratyphoid).

The baseline health data used for the estimation of health improvements are from the Global Burden of Disease 2016 (GBD 2016) for the state of Andhra Pradesh.⁶

Averted mortality is valued using two alternative methods:

- (i) Value of statistical life at 72 times GDP per capita in Andhra Pradesh, based on methodology developed by the World Bank (2016);
- (ii) Years of life lost (YLL) to premature mortality discounted at 3%, 5%, and 8% and valued at 3 times GDP per capita in Andhra Pradesh.

Reduced morbidity, or incidence of illness, is also valued using two alternative methods:

- (i) Cost-of-illness (COI) approach that includes averted health care expenditure and productivity benefits in terms of averted lost work days;

⁶ <https://vizhub.healthdata.org/gbd-compare/india>

- (ii) Years lived with disability (YLDs) valued at 3 times GDP per capita in Andhra Pradesh, where YLD is years or fraction of a year with illness or injury multiplied by a disability weight.

Productivity benefits are time savings and averted loss of work days and care giving associated with illness. Time savings are valued at 50% of average wage rates. Averted lost work days among the working population are valued at average wage rates. Averted days of care giving are valued at 50% of average wage rates. Average wage rates are estimated from GDP per capita and labor force participation rates in Andhra Pradesh, and labor income share of GDP for India. Urban and rural wage differentials are estimated from wage differentials reported in the National Sample Survey 68 (NSS 68). Male/female wage differentials are estimated from the NSS 68 and the Labor and Employment Survey 2015-16.

Table 1.1 Basic data for Andhra Pradesh

	Andhra Pradesh	Remark
Population, 2017	54,238,698	
Baseline mortality, cases in 2016	31,314	Diarrhea, intestinal infectious diseases, indirect effect on infectious disease mortality
Baseline YLDs, 2016	70,209	Diarrhea, intestinal infectious diseases
GDP per capita, 2017 Rs	123,520	
VSL to GDP per capita ratio	72	Method in World Bank and IHME (2016)
VSL, 2017 Rs	8,893,447	Product of GDP per capita and VSL to GDP per capita ratio
Average daily wage rate, 2017 Rs	466	Based on GDP per capita in 2017
Average daily wage rate, Urban, 2017 Rs	706	Based on urban/rural differentials reported in NSS 68
Average daily wage rate, Rural, 2017 Rs	353	Based on urban/rural differentials reported in NSS 68

Costs of interventions are unique to each intervention and discussed in the intervention sections.

1.3 Literature review

The main literature utilized in this paper pertains to the assessment of health benefits of interventions. Wolf et al (2014) present a meta-analysis of relative risks (RRs) of disease and mortality associated with various types of household drinking water and sanitation in low- and middle-income countries globally. Although diarrheal disease and intestinal infections are caused by many factors, the RRs provide the health benefits of improved drinking water supply and sanitation (relative to unimproved drinking water supply and unimproved or no

sanitation) as well as the health benefits of household point-of-use (POU) treatment of drinking water from various sources. The RRs associated with each intervention is applied to the baseline diarrheal and intestinal infectious disease burden to estimate health benefits. There is no need for baseline disease burden data for drinking water supply and sanitation separately.

The meta-analysis by Wolf et al include studies of 20 comparisons of alternative drinking water supplies, 48 comparisons of POU treatment vs. no POU treatment, and 12 comparisons of improved vs. unimproved sanitation. These studies were included in the meta-analysis from a review and assessment of a larger set of studies. Pruss-Ustun et al (2014) utilize the RRs in Wolf et al and provide a framework and methodology for estimating preventable disease burden from inadequate drinking water and sanitation. Olofin et al (2013) provide a methodology for estimating child mortality associated with poor nutritional status of children, and Fewtrell et al (2007) provide guidance on the magnitude of effect of diarrheal disease in early childhood on children's nutritional status (annex 1).

For improved sanitation, the RR from a study of rural households in India by Andres et al (2014) is utilized. This study assessed both the direct health benefit of improved sanitation to the household as well as the external health benefit to the community of high sanitation coverage rates. The study was also utilized by a recent report on costs and benefits of sanitation in India by UNICEF (UNICEF, 2018).

Sufficient research evidence of recent origin and high quality of the magnitude of RRs in Andhra Pradesh or even in India is not available for the other interventions. Results from the global meta-analysis by Wolf et al, as well as Olofin et al/Fewtrell et al, are therefore applied to Andhra Pradesh. This can only provide an indication, rather than a precise estimate, of the health benefits one may expect from the four interventions assessed in this paper because the health benefits depend critically on location specific factors such as natural, physical, socio-economic and behavioral factors. A sensitivity analysis of the RRs that determine the estimated magnitude of health benefits is therefore warranted. As is reported in each intervention section, the benefit-cost ratios (BCRs) of interventions are not very sensitive to changes in the RRs for some of the interventions (because of the presence of substantial non-health benefits) while for others they are.

The magnitude of health benefits of interventions is also influenced by the quality and access to public health services and medical care, as well as home management of (diarrheal) infections. These factors influence in particular case fatality rates of (diarrheal) infections. However, this is reflected in the baseline health data used for estimating the health benefits of interventions.

1.4 Household drinking water and sanitation situation

About 73% of households in Andhra Pradesh had access to an improved drinking water source in 2015-16 according to the National Family Health Survey 4 (NFHS 4). This is defined in the NFHS 4 as piped water into dwelling/yard/plot, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rainwater, and community reverse-osmosis (RO) plant.

Improved drinking water sources are generally considered having a lesser risk of pathogen contamination than unimproved sources (e.g., shallow dug wells, unprotected springs, open surface water). Water from improved sources are, however, not necessarily safe to drink (Pruss-Ustun et al, 2014). Nevertheless, only 19% of households practiced point-of-use (POU) treatment of their drinking water in 2005-06 (Andhra Pradesh and Telengana), mainly by filtering (11%) and boiling (8%) (NFHS 3). Moreover, 14% of the population in Andhra Pradesh (and Telengana) has more than 30 minutes round-trip to their drinking water source, particularly affecting women and children who often perform the task of fetching drinking water.

About 61% of households in Andhra Pradesh had a sanitary toilet facility in 2015 according to the Swachhta Status Report 2016 (MSPI/GOI, 2017). Moreover, 54% of households had access to an improved non-shared sanitation facility in 2015-16 according to the National Family Health Survey IV. Thus about 7% of households shared a sanitation facility with other households in 2015. An improved sanitation facility is defined in the NFHS 4 as flush/pour-flush toilets to piped sewer system, septic tank, or pit latrine; ventilated improved pit (VIP)/biogas latrine; pit latrine with slab; twin pit/composting toilet; which facility is not shared with any other household.

The Swachhta Status Report 2016 also reveals that 2.2% of persons with a sanitary toilet facility continued to practice open defecation (OD). Additionally, 39% of households had no sanitation facility and practiced OD in 2015 (MSPI/GOV, 2017). A substantial drive by the government for household sanitation and eradication of OD has, however, reduced this figure quite substantially since 2015.

2. Improved drinking water source

2.1 Description of intervention

About 71% of urban households in Andhra Pradesh had an improved drinking water source in 2015-16, and 74% did so in rural areas of the state (NFHS 4, 2015-16). Moreover, a substantial share of households has more than 30 minutes round-trip to their drinking water source.

The most common type of improved drinking water source is piped water to dwelling in urban areas and public tap/standpipe or tubewell/borehole in rural areas (NSS 71, 2014).

Thus the intervention is:

- i) Piped water to dwelling for urban households without an improved drinking water source.
- ii) Tubewell/borehole for rural households without an improved drinking water source or that have more than 30 minutes round-trip to their drinking water source.

Health benefits are estimated for urban and rural households that currently rely on unimproved drinking water sources. For rural households, time savings benefits are estimated for households that currently have more than 30 minutes to drinking water source, whether this source is improved or unimproved. Distant unimproved sources may not be contaminated by pathogens or other pollutants, but there is risk of pathogen contamination during extraction, transportation and storage of water prior to drinking.

2.2 Calculation of Costs and Benefits

Most of the data used for the calculation of costs and benefits of the improved drinking water intervention are presented in section 1.2. Remaining data are presented below.

2.2.1 Costs

Costs of improved drinking water supply include initial capital cost, and annual operations and maintenance (O&M) cost. No promotion program is needed as the intervention is here considered public works.

An initial capital cost of Rs. 15,000 is applied with annual O&M cost at 1.5% of initial capital cost. Annualized cost per household is estimated at Rs. 1,429 (table 2.1) and total annualized cost of intervention is estimated at Rs. 6.7 billion (table 2.2) based on total intervention beneficiaries of 4.7 million households (see next section).

Table 2.1. Cost of improved drinking water supply (Rs per household)

Capital cost	15,000
Useful life (years)	20
Annualized capital cost	1,204
O&M (1.5% of capital cost per year)	225
Total annualized intervention cost	1,429

Note: Discount rate: 5%.

Table 2.2. Total annualized cost of intervention, Rs million

	Urban	Rural	Total
Beneficiary households (000)	1,596	3,082	4,678
Total annualized cost, Rs million	2,280	4,403	6,683

Note: Discount rate: 5%. Source: Estimates by author.

2.2.2 Benefits

For the estimation of health benefits of the intervention, the population or households of Andhra Pradesh are distributed in four categories of drinking water supply according to current cover (Pre-intervention) and the expected distribution after the implementation of the intervention (Post-intervention) (table 2.3). Cover is based on NFHS 4 (2015-16) while time to drinking water source is based on projections from NFHS 3 (2005-06). While the current percent of households having more than 30 minute round-trip to drinking water source is uncertain, the pre- and post-intervention population distribution has minimal effect

on the benefit-cost ratios (BCRs) of the intervention and only on total benefits and costs of the intervention.

The risk (RR) of illness and mortality associated with the use of an improved drinking water source (i.e., the intervention) is 0.77 for urban households (piped water) and 0.89 for rural households (tubewell/borehole) *relative* to unimproved sources (see Annex 1). This means that the disease reduction from the intervention is 23% and 11% for urban and rural households respectively who currently rely on unimproved drinking water sources.

Table 2.3. Population distribution by type of drinking water supply in Andhra Pradesh

Pre-intervention	Urban	Rural	Total	RR Urban	RR Rural
Improved drinking water; access < 30 min	67%	68%	67%	0.77	0.89
Improved drinking water; access > 30 min	4%	6%	5%	0.77	0.89
Unimproved drinking water; access < 30 min	25%	20%	22%	1.0	1.0
Unimproved drinking water; access > 30 min	4%	6%	5%	1.0	1.0
Post-intervention					
Improved drinking water; access < 30 min	100%	100%	100%	0.77	0.89
Improved drinking water; access > 30 min	0%	0%	0%	0.77	0.89
Unimproved drinking water; access < 30 min	0%	0%	0%	1.0	1.0
Unimproved drinking water; access > 30 min	0%	0%	0%	1.0	1.0

Source: Author.

The intervention would benefit about 4.7 million households. The Potential Impact Fraction (PIF) formula is applied to the data in table 2.3 to estimate health benefits (see Annex 1), with estimated cases of 1,400 deaths and 3.1 million cases of diarrheal illness averted per year (table 2.4).

Table 2.4. Health benefits of intervention

	Urban	Rural	Total
Beneficiary households (000)		1,596	3,082
Averted deaths per year		765	640
Averted YLDs per year		1,714	1,434
Averted cases of diarrhea per year (000)		1,669	1,396

Source: Estimates by author.

The total annualized value of the health and productivity benefits of the intervention is estimated at Rs. 25 - 38 billion, depending on valuation method and discount rate used for valuation of deaths averted (table 2.5). The value of averted deaths accounts for 44-64% of total benefits. Health care cost savings, accounting for about 3-5% of total benefits, are

estimated based on a diarrheal disease treatment rate of 30% and a treatment cost of Rs. 1,244 per case.

Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for about 10-15% of total benefits. A loss of two work days are assumed to be averted per case of diarrhea averted among the working population, valued at average wage rates. Caregiving for children is valued at 50% of wage rates per day of child illness.

Productivity benefits in terms of time savings account for 23-36% of total benefits. Time savings are 30 minutes per household per day for those who would receive drinking water supply in the proximity of their home. These time savings are valued at 50% of female wage rates.

Table 2.5. Value of benefits of intervention, Rs million per year

	Urban	Rural	Total
Value of deaths averted			
Valuation method: VSL	13,468	11,268	24,737
Valuation method: YLL=3*GDP per capita	5,971	4,995	10,966
Health care cost savings	659	507	1,166
Productivity benefits (averted lost work days)	2,606	1,090	3,696
Productivity benefits (time savings)	3,561	5,302	8,863
Total benefits			
VSL	20,294	18,167	38,461
YLL=3*GDP per capita	12,796	11,894	24,690

Note: Discount rate: 5%. Source: Estimates by author.

An alternative valuation of health benefits is presented in table 2.6. Averted mortality is valued as in table 2.5. Reduced morbidity in terms of averted YLDs is, however, valued at 3 times GDP per capita instead of health care cost savings and averted lost work days and caregiving. This gives 7-10% lower total benefits.

Table 2.6. Alternative value of benefits of intervention, Rs million per year

	Urban	Rural	Total
Value of deaths averted			
Valuation method: VSL	13,468	11,268	24,737
Valuation method: YLL=3*GDP per capita	5,971	4,995	10,966
Value of YLD averted	1,258	1,053	2,311
Productivity benefits (time savings)	3,561	5,302	8,863
Total benefits			
VSL	18,288	17,623	35,911
YLL=3*GDP per capita	10,790	11,350	22,140

Note: Discount rate: 5%. Source: Estimates by author.

2.2.3 Benefit-cost ratios

A comparison of benefits and costs, and benefit-cost ratios (BCRs) are presented in four tables. All tables use either VSL or the value of YLL at 3 times GDP per capita for valuation of averted deaths. All tables also include the value of time savings. The two first tables use the cost-of-illness (COI) approach, i.e., health care cost savings and productivity benefits, to value the benefits of averted illness. The two last tables use valuation of YLDs as discussed above.

BCRs are larger for urban households than for rural households. This is because the percentage disease reduction from piped water supply for urban households (23%) is larger than the reduction from tubewell/borehole for rural households (11%).

BCRs using health care savings and productivity benefits are in the range of 1.9-10.7 (tables 2.7a-b). They are larger using VSL than valuation of YLL. They are also larger for urban households than for rural households.

BCRs using YLD are in the range of 1.8-9.7 (tables 2.7c-d). They are again somewhat larger using VSL than valuation of YLL, and larger for urban households than for rural households.

Table 2.7a. Benefits and costs of intervention, Rs million per year and BCRs

VSL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban	21,144	1,968	10.7	20,294	2,280	8.9	19,127	2,797	6.8
Rural	18,929	3,801	5.0	18,167	4,403	4.1	17,120	5,402	3.2

Source: Estimates by author.

Table 2.7b. Benefits and costs of intervention, Rs million per year and BCRs

YLL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban	14,939	1,968	7.6	12,796	2,280	5.6	10,788	2,797	3.9
Rural	13,738	3,801	3.6	11,894	4,403	2.7	10,143	5,402	1.9

Source: Estimates by author.

Table 2.7c. Benefits and costs of intervention, Rs million per year and BCRs

VSL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban	19,075	1,968	9.7	18,288	2,280	8.0	17,206	2,797	6.2
Rural	18,382	3,801	4.8	17,623	4,403	4.0	16,581	5,402	3.1

Source: Estimates by author.

Table 2.7d. Benefits and costs of intervention, Rs million per year and BCRs

YLL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban	12,870	1,968	6.5	10,790	2,280	4.7	8,868	2,797	3.2
Rural	13,190	3,801	3.5	11,350	4,403	2.6	9,604	5,402	1.8

Source: Estimates by author.

2.3 Assessment of Quality of Evidence

The dimensions that most importantly affect the estimated benefits and costs of the intervention are presented in table 2.8. Quantified health benefits of the intervention are proportional to the baseline health data. These data are most likely of medium-strong quality. The relative risk (RR) of disease and mortality reductions from improved drinking water supply vs. unimproved is based on a meta-analysis of 20 studies (Wolf et al, 2014). This meta-analysis found an overall RR that is quite consistent with previous meta-analyses (e.g., Waddington et al, 2009). The value of statistical life (VSL) used for valuation of mortality benefits is from a benefit-transfer function developed by the World Bank (World Bank, 2016). The function is based on meta-analysis of VSL studies from mostly high- and medium-income countries and other available evidence of VSL by country income level. Productivity benefits in terms of time savings account for about 50% of total benefits. The time savings are based on limited evidence as studies of these savings are very limited.

In terms of cost of intervention, capital cost and O&M is a rough estimate.

Table 2.8. Quality of evidence

	Quality of evidence
<i>Benefits of intervention:</i>	
Baseline health data	Medium-Strong
Relative risks (RR) for health benefits	Medium-Strong
Valuation of mortality	Medium-Strong
Productivity benefits (time savings)	Medium-Strong
<i>Cost of intervention:</i>	
Capital and O&M	Limited-Medium
Total evidence	Medium

Source: Author.

2.4 Sensitivity Analysis

An analysis is carried out as to the sensitivity of the BCRs to some of the dimensions in table 2.8: i) the magnitude of the relative risks (RRs); ii) the productivity benefits (time savings);

and iii) capital cost of intervention. The analysis assesses a 50% change in these parameters in the direction of lowering the BCRs. The sensitivity analysis is relative to the base case of using VSL for mortality and cost-of-illness (COI) for morbidity benefits, and a discount rate of 5%.

The results are presented in table 2.9:

- i) The BCR is quite sensitive to a change in the RRs in both urban and rural areas. This is because health benefits constitute such a large share of total benefits.
- ii) A change in time savings from 30 min to 15 minutes per household per day has only a small effect on the BCR. This is because of the relatively modest share of time savings in total benefits.
- iii) An increase in the capital cost of intervention from Rs. 15,000 to Rs. 22,500 reduces the BCR by one-third in both urban and rural areas.

Table 2.9 BCR sensitivity analysis

		Urban	Rural
Base case BCR		8.9	4.1
i)	BCR with 50% change in RR (urban RR=0.885; rural RR=0.945)	4.9	2.6
ii)	BCR with 50% change in time savings (15 min per household per day)	8.1	3.5
iii)	BCR with 50% higher capital cost of intervention (Rs. 22.500 per household)	5.9	2.8

Note: Discount rate: 5%. Source: Estimates by author.

3. Household point-of-use treatment of drinking water

3.1 Description of intervention

About 19% of households in Andhra Pradesh (including Telangana) practiced appropriate methods of point-of-use (POU) treatment of drinking water a decade ago, compared to over 20% nationwide in India according to the NFHS 3 (2005-06). The main appropriate methods in Andhra Pradesh were the use of a water filter (11%) and boiling (8%).

Recent data on POU treatment from the NFHS 4 (2015-16) were not publicly available at the time of writing this paper. POU treatment by appropriate methods is likely to have increased somewhat, postulated at 23% currently. Given the low rate of POU treatment by appropriate

methods, and that water filter was the most common method in Andhra Pradesh, the intervention assessed in this paper is a behavioral change campaign (BCC) that promotes household POU treatment of drinking water with water filter.

Such a BCC program needs to include an assessment of the reasons for the low rate of POU treatment of drinking water in Andhra Pradesh so that obstacles can be overcome. It also needs to raise awareness of health benefits, as well as ensure that households properly maintain the filter devices and replace necessary parts at regular intervals.

3.2 Literature Review

The challenge with an assessment of the benefits and costs of a BCC that promotes POU treatment of drinking water is the determination of the effectiveness of the campaign and the cost of the campaign in relation to its effectiveness. Three handwashing (HW) programs can shed some light on this. These programs achieved behavioral change rates of 10% to 18% of targeted households. Costs per household of the program that achieved 18% behavioral change is multiple times higher than the program that achieved 10%, demonstrating the rapidly rising marginal program cost in relation to behavioral change rates (table 3.1).

“Low”, “mid” and “high” intensity promotion programs of 10%, 15% and 20% behavioral change rate is applied in this paper to the BCC for promotion of POU treatment of drinking water, with costs in table 3.1 adjusted for annual inflation to 2017.

Table 3.1. Cost of behavioral change promotion

	Guatemala	Thailand	Burkina Faso
Behavioral change (% of target population)	10%	16%	18%
Program Cost per Target Household (US \$)	0.35	1.2	5.0
Program Cost per Household with Behavioral Change (US \$)	3.6	7.4	27.9

Source: Derived from Saade et al (2001), Pinfold and Horan (1996), and Borghi et al (2002).

3.3 Calculation of Costs and Benefits

Most of the data used for the calculation of costs and benefits of the BCC intervention for POU treatment are presented in section 1.2. Remaining data are presented below.

3.3.1 Costs

An average water filter for POU of drinking water costs about Rs. 2,000 based on an online market review.⁷ Parts replacement may be around Rs. 500 per year. There is also a cost of time spent on filtering water, here 2 minutes per household per day valued at 50% of wage rates. These costs are applicable to around 1.4-2.9 million households expected to start POU of drinking water as a result of BCC promotion program (table 3.2).

Table 3.2. Cost of POU treatment of drinking water

	Low	Mid	High
Households starting POU of drinking water (000)	1,429	2,143	2,858
Cost of water filter (Rs per household)	2000	2000	2000
Useful life (years)	5	5	5
Annualized capital cost (Rs per household)	462	462	462
Parts replacement (1 time per year), Rs per household	500	500	500
Cost of time spent on water filtering (Rs/household/year)	434	434	434
Sub-total annualized cost, Rs million (all households)	1,995	2,993	3,991

Note: Discount rate: 5%. Source: Author.

The BCC program promoting POU treatment of drinking water is undertaken once over the assumed 5 years of useful life of the water filter. Costs are according to table 3.1 adjusted for annual inflation. The program is directed at all households (table 3.3). The cost of the BCC program is only substantial in comparison with the private costs in table 3.2 for the “high” intensity program.

Table 3.3. Cost of BCC program

	Low	Mid	High
Program promotion cost per household (US\$)	0.35	1.20	5.00
Program promotion cost per household (US\$), 2017	0.5	2.0	7.4
Program promotion cost per household (Rs)	30	130	480
Annualized program promotion cost per HH (Rs)	6.9	30	111
Households targeted by POU promotion (000)	14,289	14,289	14,289
Total annualized program promotion cost (Rs million)	99	429	1,584

Note: Discount rate: 5%. Source: Author.

5.3.2 Benefits

For the estimation of health benefits of the intervention, the population or households of Andhra Pradesh are distributed in two categories of household currently practicing and not practicing POU treatment of drinking water (Pre-intervention) and the expected distribution after the implementation of the intervention (Post-intervention) (table 3.4). While the

⁷ <https://www.amazon.in/Water-Filters-Purifiers/b?ie=UTF8&node=1380262031>

current POU treatment rate is uncertain, the pre-intervention population distribution has minimal effect on the benefit-cost ratios (BCRs) of the intervention and only on total benefits and costs of the intervention.

The relative risks (RRs) of illness and mortality are weighted average RR for POU treatment of piped drinking water in urban and other improved drinking water sources in rural (Pruss-Ustun et al, 2014) . The RR means that the disease reduction from POU treatment is 35%. The BCC intervention effectiveness is discussed above, ranging from 10% to 20%.

Table 3.4. Population distribution by household POU treatment of drinking water in Andhra Pradesh

	Low	Mid	High	RR
Pre-intervention				
POU treatment (appropriate method)	23%	23%	23%	0.65
Not appropriate or no treatment	77%	77%	77%	1.0
BBC Water Filter promotion effectiveness (% behavioral change rate)	10%	15%	20%	
Post-intervention				
POU treatment (appropriate method)	33%	38%	43%	0.65
Not appropriate or no treatment	67%	62%	57%	1.0

Source: Author.

The intervention is expected to make 1.4-2.9 million households start POU treatment of drinking water. The Potential Impact Fraction (PIF) formula is applied to the data in table 3.4 to estimate health benefits (see Annex 1), with estimated cases of about 1,200-2,400 deaths and 2.6-5.1 million cases of diarrheal illness averted per year (table 3.5).

Table 3.5. Health benefits of intervention

	Low	Mid	High
Households starting POU of drinking water (000)	1,429	2,143	2,858
Averted deaths per year	1,180	1,769	2,359
Averted YLDs per year	2,645	3,967	5,290
Averted cases of diarrhea per year (000)	2,574	3,861	5,149

Source: Estimates by author.

The total annualized value of the health and productivity benefits of the intervention is estimated at Rs. 8 – 31 billion (table 3.6). The value of averted deaths accounts for 69-83% of total benefits. Health care cost savings, accounting for about 6-11% of total benefits, are estimated based on a diarrheal disease treatment rate of 30% and a treatment cost of Rs. 1,244 per case.

Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for somewhat 11-20% of total benefits. A loss of two work days are assumed to be averted per case of diarrhea averted among the working population, valued at average wage rates. Caregiving for children is valued at 50% of wage rates per day of child illness.

Table 3.6. Value of benefits of intervention, Rs million per year

	Low	Medium	High
Value of deaths averted			
VSL	12,863	19,295	25,727
YLL=3*GDP per capita	5,702	8,553	11,405
Health care cost savings	919	1,379	1,838
Productivity benefits (averted lost work days and caregiving)	1,642	2,463	3,284
Total benefits			
VSL	15,424	23,136	30,848
YLL=3*GDP per capita	8,263	12,395	16,526

Note: Discount rate: 5%. Source: Estimates by author.

An alternative valuation of health benefits is presented in table 3.7. Averted mortality is valued as in table 3.6. Reduced morbidity in terms of averted YLDs is, however, valued at 3 times GDP per capita instead of health care cost savings and averted lost work days and caregiving. This gives 9-17% lower total benefits.

Table 3.7. Alternative value of benefits of intervention, Rs million per year

	Low	Medium	High
Value of deaths averted			
VSL	12,863	19,295	25,727
YLL=3*GDP per capita	5,702	8,553	11,405
Value of YLD averted	1,202	1,803	2,403
Total benefits			
VSL	14,065	21,097	28,130
YLL=3*GDP per capita	6,904	10,356	13,808

Note: Discount rate: 5%. Source: Estimates by author.

3.3.3 Benefit-cost ratios

A comparison of benefits and costs, and benefit-cost ratios (BCRs) are presented in four tables, as for the previous interventions. All tables use either VSL or the value of YLL at 3 times GDP per capita for valuation of averted deaths. The two first tables use the cost-of-illness (COI) approach, i.e., health care cost savings and productivity benefits, to value the benefits of averted illness. The two last tables use valuation of YLDs as discussed above.

BCRs using health care savings and productivity benefits are in the range of 2.4-7.5 (tables 3.8a-b). They are larger using VSL than valuation of YLL.

BCRs using YLD are in the range of 1.9-6.9 (tables 3.8c-d). They are again larger using VSL than valuation of YLL.

The BCRs are declining with increased intensity (from “low” to “high”) of the BCC program, because of rising unit costs of achieving behavioral change.

Table 3.8a. Benefits and costs of intervention, Rs million per year and BCRs

VSL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	15,464	2,055	7.5	15,424	2,094	7.4	15,367	2,156	7.1
Mid	23,195	3,347	6.9	23,136	3,422	6.8	23,050	3,538	6.5
High	30,927	5,420	5.7	30,848	5,575	5.5	30,734	5,815	5.3

Source: Estimates by author.

Table 3.8b. Benefits and costs of intervention, Rs million per year and BCRs

YLL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	9,766	2,055	4.8	8,263	2,094	3.9	6,936	2,156	3.2
Mid	14,649	3,347	4.4	12,395	3,422	3.6	10,403	3,538	2.9
High	19,533	5,420	3.6	16,526	5,575	3.0	13,871	5,815	2.4

Source: Estimates by author.

Table 3.8c. Benefits and costs of intervention, Rs million per year and BCRs

VSL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	14,103	2,055	6.9	14,065	2,094	6.7	14,010	2,156	6.5
Mid	21,154	3,347	6.3	21,097	3,422	6.2	21,014	3,538	5.9
High	28,206	5,420	5.2	28,130	5,575	5.0	28,019	5,815	4.8

Source: Estimates by author.

Table 3.8d. Benefits and costs of intervention, Rs million per year and BCRs

YLL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	8,406	2,055	4.1	6,904	2,094	3.3	5,578	2,156	2.6
Mid	12,608	3,347	3.8	10,356	3,422	3.0	8,367	3,538	2.4
High	16,811	5,420	3.1	13,808	5,575	2.5	11,157	5,815	1.9

Source: Estimates by author.

3.4 Assessment of Quality of Evidence

The dimensions that most importantly affect the estimated benefits and costs of the intervention are presented in table 3.9. The quality of evidence of the baseline health data

and valuation of mortality is discussed before. The relative risk (RR) of disease and mortality reductions from POU treatment of drinking water is based on a meta-analysis of 48 studies (Wolf et al, 2014).

In terms of cost of intervention, the cost of water filter and maintenance is by far the largest cost component and has medium-strong evidence.

Table 3.9. Quality of evidence

	Quality of evidence
<i>Benefits of intervention:</i>	
Baseline health data	Medium-Strong
Relative risks (RR) for health benefits	Medium-Strong
Valuation of mortality	Medium-Strong
<i>Cost of intervention:</i>	
Cost of water filter and maintenance	Medium-Strong
<i>Total evidence</i>	Medium-Strong

Source: Author.

3.5 Sensitivity Analysis

An analysis is carried out as to the sensitivity of the BCRs to two of the dimensions in table 3.9: i) the magnitude of the relative risks (RRs); and ii) cost of intervention. The analysis assesses a 50% change in these parameters in the direction of lowering the BCRs. The sensitivity analysis is relative to the base case of using VSL for mortality and cost-of-illness (COI) for morbidity benefits, and a discount rate of 5%.

The results are presented in table 3.10:

- i) The sensitivity of the BCR is proportional to the change in RR. This is because the benefits are entirely health benefits.
- ii) An increase of 50% in each of the components of the cost of intervention separately reduces the BCR only somewhat. A simultaneous increase in the cost of all components reduces the BCR by one-third.

Table 3.10. BCR sensitivity analysis

		Mid-case
Base case BCR		6.8
i)	BCR with 50% change in RR (RR=0.865)	3.3
ii)	BCR with 50% higher cost of water filter (Rs. 3,000 per household)	5.9
	BCR with 50% higher cost of parts replacement (Rs. 750 per year per household)	5.8
	BCR with 50% higher cost of time spent on treatment (3 minutes per household per day)	6.0
	BCR with 50% higher program promotion cost	6.4
	BCR with 50% higher cost of all cost components	4.5

Source: Estimates by author.

4. Improved sanitation

4.1 Description of intervention

About 61% of households in Andhra Pradesh had a sanitary toilet facility in 2015 according to the Swachhta Status Report 2016 (MSPI/GOI, 2017). Moreover, 54% of households had access to an improved non-shared sanitation facility in 2015-16 according to the National Family Health Survey IV.⁸ Thus about 7% of households shared a sanitation facility with other households in 2015. An improved non-shared sanitation facility is defined in the NFHS 4 as flush/pour-flush toilets to piped sewer system, septic tank, or pit latrine; ventilated improved pit (VIP)/biogas latrine; pit latrine with slab; twin pit/composting toilet; which facility is not shared with any other household.

The Swachhta Status Report 2016 also reveals that 2.2% of persons with a sanitary toilet facility continued to practice open defecation (OD). Additionally, 39% of households had no sanitation facility and practiced OD in 2015 (MSPI/GOV, 2017). A substantial drive by the government for household sanitation and eradication of OD has, however, reduced this figure quite substantially since 2015. Nevertheless, sanitation programs inevitably encounter challenges of quality of construction, choice of appropriate sanitation technology, and achieving sustained OD free communities.

⁸ The Joint Monitoring Programme (JMP) of WHO/UNICEF classifies household sanitation into improved and unimproved facilities.

The most common household sanitation facility in Andhra Pradesh is a flush/pour-flush toilet in both urban and rural areas. The Government of India is promoting household sanitation facilities with a monetary incentive up to Rs 12,000 to eligible households. Most households opt for a flush/pour-flush system with a single- or twin-pit. This is therefore the intervention assessed in this paper, with the aim of reaching 95% household coverage with improved, non-shared sanitation. The government may, however, aim for 100% coverage. The difference has no/negligible effect on the benefit-cost ratios (BCRs), unless the cost of reaching the last 5% of households escalates.

Institutional sanitation (sanitation facilities at pre-schools, schools, health facilities, etc.), albeit important, is not assessed in this paper.

4.2 Calculation of Costs and Benefits

Most of the data used for the calculation of costs and benefits of the sanitation intervention are presented in section 1.2. Remaining data are presented below.

4.2.1 Costs

Costs of household sanitation include initial capital cost, periodic emptying of pit or septic tank, regular cleaning of the facility, annual operations and maintenance (O&M) cost, and cost of sanitation promotion programs by the government.

An initial capital cost of Rs. 20,000 is applied. This is for the toilet, installation, and construction of superstructure (base, building). This is the cost reported in a recent survey of over 10,000 households in 12 states conducted by UNICEF in 2017 (UNICEF, 2018), reflecting the cost paid by households and cost paid by the government among the poorest two quintiles of households, which are the households that are most likely to still not have improved sanitation. The cost may be higher or lower for some households. The impact of this on the benefit-cost ratios (BCR) is assessed in the sensitivity analysis.

Each of the government recommended twin-pits is 4-feet deep and 3-4 feet wide and takes an average household about 5 years to fill. Once one is full, the second pit is put into use

while the fecal matter in the first pit breaks down. Emptying the pit costs around Rs 1,500.⁹ Cleaning of the sanitation facility is assumed to take 10 minutes per day, with time valued at 50% of urban and rural wage rates.¹⁰ Sanitation program cost of Rs. 600 per household (5% of government sanitation budget) that currently do not have improved non-shared sanitation is assumed to cover promotion of household sanitation, proper designs, and promotion of household members full use of the facility.

Annualized cost per rural and urban household is estimated at Rs. 5,577 – 8,229 (table 4.1) and total annualized cost of intervention is estimated at Rs. 32 billion (table 4.2) based on total intervention beneficiaries of 5.4 million households (see next section). Cost of cleaning constitutes over half of the cost for urban households and nearly half of the cost for rural households.

Table 4.1. Cost of household sanitation (Rs per household)

	Urban	Rural	Remark
Capital cost	20,000	20,000	Initial cost
Emptying of pit	1,500	1,500	Once per 5 years
Cost of cleaning per year	5,304	2,652	10 minutes per household per day
O&M cost	1,000	1,000	5% of capital cost per year
Program cost	600	600	Once per household
Annualized cost, Rs	8,229	5,577	Per household

Note: Discount rate: 5%. Source: The author.

Table 4.2. Total annualized cost of intervention, Rs million

	Urban	Rural	Total
Beneficiary households (000)	661	4,785	5446
Total annualized cost, Rs million	5,441	26,683	32,124

Note: Discount rate: 5%. Source: Estimates by author.

4.2.2 Benefits

For the estimation of health benefits of the intervention, the population or households of Andhra Pradesh are distributed in four categories of household sanitation according to current sanitation cover (Pre-intervention) and the expected distribution after the implementation of the intervention which aims to reduce the population with shared and unimproved or no sanitation to 5% in both urban and rural areas (Post-intervention) (table

⁹ Useful sanitation information from Rajasthan at <https://scroll.in/article/759201/swachh-bharat-in-Rajasthan-a-village-shows-why-open-defecation-is-ending-only-on-paper>

¹⁰ 10 minutes per day is here considered sufficient in order to reap the diarrheal risk reduction applied in this paper. Households may, however, spend more time on cleaning for aesthetic and status benefits that are not captured in this paper. In the survey by UNICEF in July-August 2017, households report an average cleaning time of the sanitation facility of about 30 minutes per day.

4.3). Cover is based on the Swachhta Status Report 2016 (MSPI/GOI, 2017) and NFHS 4 (2015-16) for Andhra Pradesh. Non-use of existing facilities is based on the Swachhta Status Report 2016. While the current sanitation cover and non-use of existing facilities is uncertain in light of the government's drive to increase coverage, the pre- and post-intervention population distribution has minimal effect on the benefit-cost ratios (BCRs) of the intervention and only on total benefits and costs of the intervention.

The risk (RR) of illness and mortality associated with the use of an improved, non-shared sanitation facility (i.e., the intervention) is 0.72 *relative* to unimproved or no sanitation facility. This includes the direct health benefit to the household adopting improved sanitation and the external health benefit to the community, as estimated by Andres et al (2014) (see Annex 1). The RR associated with the use of shared sanitation facilities is here assumed to be 0.77. Thus disease reduction from the intervention is 47% and 23%, respectively.

Table 4.3. Population distribution by type of household sanitation in Andhra Pradesh

Pre-intervention	Urban	Rural	Total	RR
Have improved non-shared sanitation and using it	76.4%	41.8%	52.3%	0.53
Have improved non-shared sanitation BUT not using it	1.0%	1.3%	1.3%	1.0
Shared sanitation	11.4%	4.8%	7.4%	0.77
Other unimproved or no sanitation	11.2%	52.1%	39.0%	1.0
Post-intervention				
Intervention target (access rate of improved non-shared sanitation)	95%	95%	95%	
Have improved non-shared sanitation and using it	94.0%	92.4%	92.9%	0.53
Have improved non-shared sanitation BUT not using it	1.0%	2.6%	2.1%	1.0
Shared sanitation	0.6%	0.2%	0.4%	0.77
Other unimproved on no sanitation	4.4%	4.8%	4.6%	1.0

Source: Author.

The intervention would benefit about 5.4 million households. The Potential Impact Fraction (PIF) formula is applied to the data in table 4.3 to estimate health benefits (see Annex 1), with estimated cases of over 7,000 deaths and 15.4 million cases of diarrheal illness averted per year (table 4.4).

Table 4.4. Health benefits of intervention

	Urban	Rural	Total
Beneficiary households (000)	661	4,785	5,446
Averted deaths per year	928	6,113	7,041
Averted YLDs per year	2,080	13,706	15,786
Averted cases of diarrhea per year (000)	2,024	13,341	15,365

Source: Estimates by author.

The total value of the health and productivity benefits of the intervention is estimated at Rs. 172 – 241 billion, depending on valuation method and discount rate used for valuation of deaths averted (table 4.5). The value of averted deaths accounts for 32-52% of total benefits. Health care cost savings, accounting for 2-3% of total benefits, are estimated based on a diarrheal disease treatment rate of 30% and a treatment cost of Rs. 1,244 per case.

Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for 6-8% of total benefits. A loss of two work days are assumed to be averted per case of diarrhea averted among the working population, valued at average wage rates. Caregiving for children is valued at 50% of wage rates per day of child illness.

Productivity benefits in terms of time savings account for 4-57% of total benefits. Time savings are 20 minutes per household member per day for those who no longer would practice open defecation (OD) and 3 minutes per household member per day for those no longer with a shared sanitation facility. These time savings are valued at 50% of wage rates for all beneficiaries 5+ years of age. In the recent survey by UNICEF household members reported that they on average spend 40 minutes per day on OD (UNICEF, 2018). This may, however, include time spent on co-activities that household members may undertake. Thus, to be conservative, only 20 minutes are accounted as time savings of ending OD in this paper. This is also the average time spent on OD reported by households in 6 East Asian countries according to the Economics of Sanitation Initiative (ESI) by the Water and Sanitation Program, World Bank.¹¹

Table 4.5. Value of benefits of intervention, Rs million per year

	Urban	Rural	Total
Deaths averted			
VSL	16,339	107,673	124,012
YLL=3*GDP per capita	7,243	47,732	54,975
Health care cost savings	799	4,841	5,640
Productivity benefits (averted lost work days)	3,161	10,417	13,578
Productivity benefits (time savings)	13,227	84,220	97,447
Total benefits, Rs million			
VSL	33,527	207,151	240,678
YLL=3*GDP per capita	24,431	147,210	171,640

Note: Discount rate: 5%. Source: Estimates by author.

¹¹ <https://www.wsp.org/content/economic-impacts-sanitation>

An alternative valuation of health benefits is presented in table 4.6. Averted mortality is valued as in table 4.5. Reduced morbidity in terms of averted YLDs is, however, valued at 3 times GDP per capita instead of health care cost savings and averted lost work days and caregiving. This gives slightly lower total benefits.

Table 4.6. Alternative value of benefits of intervention, Rs million per year

	Urban	Rural	Total
Deaths averted			
VSL	16,339	107,673	124,012
YLL=3*GDP per capita	7,243	47,732	54,975
Value of YLD averted	1,526	10,059	11,585
Productivity benefits (time savings)	13,227	84,220	97,447
Total benefits, Rs million			
VSL	31,092	201,953	233,045
YLL=3*GDP per capita	21,996	142,011	164,007

Note: Discount rate: 5%. Source: Estimates by author.

4.2.3 Benefit-cost ratios

A comparison of benefits and costs, and benefit-cost ratios (BCRs) are presented in four tables. All tables use either VSL or the value of YLL at 3 times GDP per capita for valuation of averted deaths. All tables also include the value of time savings. The two first tables use the cost-of-illness (COI) approach, i.e., health care cost savings and productivity benefits, to value the benefits of averted illness. The two last tables use valuation of YLDs as discussed above.

Benefits and costs are larger in rural than urban areas as most of the beneficiaries are rural. BCRs using health care savings and productivity benefits are in the range of 3.9-8.3 (tables 4.7a-b). They are somewhat larger using VSL than valuation of YLL. They are also larger for rural households than for urban households.

BCRs using YLD are in the range of 3.5-8.1 (tables 4.7c-d). They are again somewhat larger using VSL than valuation of YLL, and larger for rural households than for urban households.

Table 4.7a. Benefits and costs of intervention, Rs million per year and BCRs

VSL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban sanitation	34,941	5,422	6.4	33,527	5,441	6.2	31,584	5,518	5.7
Rural sanitation	215,880	25,998	8.3	207,151	26,683	7.8	195,164	27,987	7.0

Source: Estimates by author.

Table 4.7b. Benefits and costs of intervention, Rs million per year and BCRs

YLL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban sanitation	27,413	5,422	5.1	24,431	5,441	4.5	21,468	5,518	3.9
Rural sanitation	166,271	25,998	6.4	147,210	26,683	5.5	128,500	27,987	4.6

Source: Estimates by author.

Table 4.7c. Benefits and costs of intervention, Rs million per year and BCRs

VSL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban sanitation	32,431	5,422	6.0	31,092	5,441	5.7	29,254	5,518	5.3
Rural sanitation	210,647	25,998	8.1	201,953	26,683	7.6	190,012	27,987	6.8

Source: Estimates by author.

Table 4.7d. Benefits and costs of intervention, Rs million per year and BCRs

YLL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Urban sanitation	24,903	5,422	4.6	21,996	5,441	4.0	19,138	5,518	3.5
Rural sanitation	161,038	25,998	6.2	142,011	26,683	5.3	123,347	27,987	4.4

Source: Estimates by author.

4.3 Assessment of Quality of Evidence

The dimensions that most importantly affect the estimated benefits and costs of the intervention are presented in table 4.8. The quality of evidence of the baseline health data and valuation of mortality is discussed before. The relative risk (RR) of disease and mortality reductions from improved sanitation vs. unimproved sanitation is based on one study of Indian households. The time savings from ending OD are conservative compared to UNICEF (2018) and similar to evidence from other Asian countries.

In terms of cost of intervention, capital cost is based on recent experience of the government sanitation promotion program. Time needed on cleaning of sanitation facility to reap the health benefits estimated in this paper is uncertain.

Table 4.8. Quality of evidence

	Quality of evidence
<i>Benefits of intervention:</i>	
Baseline health data	Medium-Strong
Relative risks (RR) for health benefits	Medium
Valuation of mortality	Medium-Strong
Productivity benefits (time savings)	Limited
<i>Cost of intervention:</i>	
Capital cost	Medium-Strong
Cleaning of sanitation facility	Medium
Total evidence	Medium-Strong

Source: Author.

4.4 Sensitivity Analysis

An analysis is carried out as to the sensitivity of the BCRs to some of the dimensions in table 4.8: i) the magnitude of the relative risks (RRs); ii) the productivity benefits (time savings); and iii) cost of two intervention cost components. The analysis assesses a 50% change in these parameters in the direction of lowering the BCRs. The sensitivity analysis is relative to the base case of using VSL for mortality and cost-of-illness (COI) for morbidity benefits, and a discount rate of 5%.

The results are presented in table 4.9:

- i) The BCRs decline with 28-35% with the change in RR. This relatively large change in the BCRs is because health benefits are a major share of total benefits.
- ii) A change in time savings from 20 min to 10 minutes per person per day decreases the BCRs by no more than 18-21%.
- iii) An increase in the capital cost of intervention from Rs. 20,000 to Rs. 30,000 reduces the BCR by 15-20%. A higher cleaning cost of 20 minutes per day reduces the BCRs by 32-40% due to the high cost of cleaning relative to total cost.

A sensitivity analysis is also undertaken with respect to sanitation coverage rates. If the rural population with a sanitary toilet is 70-80% today instead of 48% in 2015 (i.e., improved non-shared and shared in table 4.3), the BCR changes minimally. Thus the actual coverage rate has no significant effect on the BCR.

Table 4.9 BCR sensitivity analysis

		Urban	Rural
Base case BCR		6.2	7.8
i)	BCR with 50% change in RR (RRs=0.76 and 0.88)	3.9	5.2
ii)	BCR with 50% change in time savings (10 min per person per day)	5.1	6.2
iii)	BCR with 50% higher capital cost of intervention (Rs. 30,000 per household)	5.3	6.3
	BCR with 50% higher cleaning cost (20 minutes per day)	3.7	5.3

Source: Estimates by author.

5. Promotion of the use of sanitation facilities

5.1 Description of intervention

The construction of a household sanitation facility does not necessarily mean that all household members will use the new facility. The Swachhta Status Report 2016 found that 2.2% of household members continue to practice OD after construction of sanitation facility. Non-use is more prevalent in rural areas (2.7%) than in urban areas (1.1%).

Faced with a situation of OD among households with sanitation facility, the intervention is a behavioral change campaign (BCC) that promotes the consistent use of existing sanitation facilities.

5.2 Literature Review

The challenge with an assessment of the benefits and costs of a BCC that promotes the use of sanitation is the determination of the effectiveness of the campaign and the cost of the campaign in relation to its effectiveness. Three handwashing (HW) programs can shed some light on this, and are reported in the section on the POU treatment of drinking water intervention.

“Low”, “mid” and “high” intensity promotion programs of 10%, 15% and 20% behavioral change rate is applied in this paper to the BCC for promotion of the use of sanitation, with costs in table 5.1 adjusted for annual inflation to 2017.

5.3 Calculation of Costs and Benefits

Most of the data used for the calculation of costs and benefits of the BCC intervention for the use of sanitation facilities are presented in section 1.2. Remaining data are presented below.

5.3.1 Costs

The BCC program promoting the use of existing household sanitation is repeated every 3 years over the useful life of the sanitation facilities. Annualized cost per household ranges from Rs. 11 for the “low” intensity program to Rs. 181 for the “high” intensity. The program is directed at all households that currently have improved sanitation, as precisely which

households have members who do not utilize existing facilities is unknown.. Thus total annualized cost is in the range of Rs. 99-1,580 million (table 5.1).

Table 5.1. Cost of BCC program promotion intervention

	Low	Mid	High
Useful life of existing sanitation facility (years)	20	20	20
BBC program promotion cost per household (US\$)	0.35	1.20	5.00
Program promotion cost per household (US\$), 2017	0.5	2.0	7.4
Program promotion cost per household (Rs)	30	130	480
Annualized program promotion cost per household (Rs)	11	49	181
Households targeted by BCC program intervention (million)	8.7	8.7	8.7
Total annualized BCC program promotion cost, Rs Million	99	428	1,580

Note: Discount rate is 5%. BBC program is repeated every 3 years over the useful life of the sanitation facility.

Source: Author.

The NSS 69 (2012) finds several reasons for non-use of existing sanitation facilities: i) no toilet superstructure; ii) lack of cleanliness/insufficient water; iii) malfunctioning of the latrine; iv) personal preferences; and v) other reasons. The two most important reasons in Andhra Pradesh were personal preferences (17%) and lack of cleanliness/insufficient water (62%).

Personal preferences for OD imply that these household members perceive some benefits of continuing OD, or cost of giving up their old practice of OD. The magnitude of this cost is unknown. It is therefore in this paper assumed to be (at least) as large as the value of time spent on OD, i.e., 20 minutes per person per day valued at 50% of average wages. This is applied to the 17% of non-users who state “personal preferences” as reason for not using existing toilet facilities (annualized cost of ending OD in table 5.2).

Moreover, that lack of cleanliness/insufficient water was reported as the second main reason for not using the toilet facility prompts the inclusion of cost of cleaning of toilet facility (or value of time spent on cleaning), i.e., 20 minutes per week valued at 50% of wages. This cost is applied to all households with non-users of existing toilet facilities (annual cost of cleaning toilets in table 5.2).

Total private cost of intervention is therefore estimated at Rs. 164 – 328 million, depending on intensity or level of effort of the promotion program and household member response rates. Total private and public (promotion program) cost is therefore Rs. 263 – 1,908.

Table 5.2. Addition cost of intervention

	Low	Mid	High
Annualized cost (foregone benefits) of ending OD	88	131	175
Annual cost of cleaning toilets	76	114	153
Total annual cost	164	245	328

Note: Discount rate: 5%. Source: Author.

5.3.2 Benefits

For the estimation of health benefits of the intervention, the population or households of Andhra Pradesh are distributed in four categories of household sanitation according to current sanitation cover and use of sanitation (Pre-intervention) and the expected distribution after the implementation of the intervention which aims to reduce the population not using existing sanitation facilities (Post-intervention) (table 5.3). The pre-intervention distribution is the same as for the improved sanitation intervention in the previous section. And again, while the current sanitation cover and non-use of existing facilities is uncertain, the pre-intervention population distribution has minimal effect on the benefit-cost ratios (BCRs) of the intervention and only on total benefits and costs of the intervention.

The risk (RR) of illness and mortality associated with the use of an improved, non-shared sanitation facility (i.e., the intervention) is 0.72 relative to unimproved or no sanitation facility (Wolf et al, 2014) (see Annex 1)¹². The RR associated with the use of shared sanitation facilities is not reported in Pruss-Ustun et al (2014) or Wolf et al (2014) and is here assumed to be 0.86. Thus disease reduction from the intervention is 28% and 14%, respectively. The BCC intervention effectiveness is discussed above, ranging from 10% to 20%.

¹² This intervention involves a relatively small number of households. The community health benefit estimated in Andres et al (2014) is therefore likely to be minimal. The relative risk in Wolf et al (2014) is therefore applied instead, reflecting primarily the direct health benefit to the household.

Table 5.3. Population distribution by type of household sanitation in Andhra Pradesh

	Low	Mid	High	RR
Pre-intervention				
Have improved non-shared sanitation and using it	52.3%	52.3%	52.3%	0.72
Have improved non-shared sanitation BUT not using it	1.3%	1.3%	1.3%	1.0
Shared sanitation	7.4%	7.4%	7.4%	0.86
Other unimproved on no sanitation	39.0%	39.0%	39.0%	1.0
BCC intervention effectiveness (% behavioral change rate)	10%	15%	20%	
Post-intervention				
Have improved non-shared sanitation and using it	52.4%	52.5%	52.5%	0.72
Have improved non-shared sanitation BUT not using it	1.2%	1.1%	1.1%	1.0
Shared sanitation	7.4%	7.4%	7.4%	0.86
Other unimproved on no sanitation	39.0%	39.0%	39.0%	1.0

Source: Author.

The intervention would end OD among 72-145 thousand household members. The Potential Impact Fraction (PIF) formula is applied to the data in table 5.3 to estimate health benefits (see Annex 1), with estimated cases of 14-28 deaths and about 30-61 thousand cases of diarrheal illness averted per year (table 5.4).

Table 5.4. Health benefits of intervention

	Low	Mid	High
Individuals ending open defecation (000)	72	109	145
Averted deaths per year	14	21	28
Averted YLDs per year	31	47	62
Averted cases of diarrhea per year (000)	30	45	61

Source: Estimates by author.

The total annualized value of the health and productivity benefits of the intervention is estimated at Rs. 0.7 – 1.6 billion (table 5.5). The value of averted deaths accounts for 16-31% of total benefits. Health care cost savings, accounting for less than 2% of total benefits, are estimated based on a diarrheal disease treatment rate of 30% and a treatment cost of Rs. 1,244 per case.

Productivity benefits in terms of averted lost work days (including caregiving by adults for ill children) account for about 4-5% of total benefits. A loss of two work days are assumed to be averted per case of diarrhea averted among the working population, valued at average wage rates. Caregiving for children is valued at 50% of wage rates per day of child illness.

Productivity benefits in terms of time savings account for 64-77% of total benefits. Time savings are 20 minutes per household member per day for those who no longer would

practice open defecation (OD), as in the case of the previous sanitation intervention. These time savings are valued at 50% of wage rates for all beneficiaries regardless of age.

Table 5.5. Value of benefits of intervention, Rs million per year

	Low	Medium	High
Value of deaths averted			
VSL	244	367	489
YLL=3*GDP per capita	108	162	217
Health care cost savings	11	17	23
Productivity benefits (averted lost work days)	31	47	62
Productivity benefits (time savings)	508	762	1,016
Total benefits, Rs million			
VSL	795	1,192	1,590
YLL=3*GDP per capita	659	988	1,317

Note: Discount rate: 5%. Source: Estimates by author.

An alternative valuation of health benefits is presented in table 5.6. Averted mortality is valued as in table 5.5. Reduced morbidity in terms of averted YLDs is, however, valued at 3 times GDP per capita instead of health care cost savings and averted lost work days and caregiving. This gives 3% lower total benefits.

Table 5.6. Alternative value of benefits of intervention, Rs million per year

	Low	Medium	High
Value of deaths averted			
VSL	244	367	489
YLL=3*GDP per capita	108	162	217
Value of YLD averted	23	34	46
Productivity benefits (time savings)	508	762	1,016
Total benefits, Rs million			
VSL	775	1,163	1,550
YLL=3*GDP per capita	639	959	1,278

Note: Discount rate: 5%. Source: Estimates by author.

5.3.3 Benefit-cost ratios

A comparison of benefits and costs, and benefit-cost ratios (BCRs) are presented in four tables, as for the previous interventions. All tables use either VSL or the value of YLL at 3 times GDP per capita for valuation of averted deaths. All tables also include the value of time savings. The two first tables use the cost-of-illness (COI) approach, i.e., health care savings and productivity benefits, to value the benefits of averted illness. The two last tables use valuation of YLDs as discussed above.

BCRs using health care savings and productivity benefits are in the range of 0.6-3.1 (tables 5.7a-b). They are somewhat larger using VSL than valuation of YLL.

BCRs using YLD are in the range of 0.6-3.0 (tables 5.7c-d). They are again somewhat larger using VSL than valuation of YLL.

Table 5.7a. Benefits and costs of intervention, Rs million per year and BCRs

VSL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	829	267	3.11	795	262	3.03	748	258	2.91
Mid	1,243	671	1.85	1,192	673	1.77	1,123	679	1.65
High	1,657	1,873	0.88	1,590	1,907	0.83	1,497	1,963	0.76

Source: Estimates by author.

Table 5.7b. Benefits and costs of intervention, Rs million per year and BCRs

YLL + health care and productivity	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	716	267	2.69	659	262	2.51	597	258	2.32
Mid	1,074	671	1.60	988	673	1.47	896	679	1.32
High	1,432	1,873	0.76	1,317	1,907	0.69	1,194	1,963	0.61

Source: Estimates by author.

Table 5.7c. Benefits and costs of intervention, Rs million per year and BCRs

VSL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	808	267	3.03	775	262	2.95	729	258	2.83
Mid	1,213	671	1.81	1,163	673	1.73	1,094	679	1.61
High	1,617	1,873	0.86	1,550	1,907	0.81	1,459	1,963	0.74

Source: Estimates by author.

Table 5.7d. Benefits and costs of intervention, Rs million per year and BCRs

YLL + YLD	3% discount rate			5% discount rate			8% discount rate		
	Benefit	Cost	BCR	Benefit	Cost	BCR	Benefit	Cost	BCR
Low	696	267	2.61	639	262	2.43	578	258	2.24
Mid	1,044	671	1.56	959	673	1.42	867	679	1.28
High	1,392	1,873	0.74	1,278	1,907	0.67	1,156	1,963	0.59

Source: Estimates by author.

5.4 Assessment of Quality of Evidence

The dimensions that most importantly affect the estimated benefits and costs of the intervention are presented in table 5.8. The quality of evidence of the benefits is discussed before.

In terms of cost of intervention, the cost of ending OD (foregone benefit) is very uncertain with limited evidence. As this is the key parameter driving the BCRs, the overall quality of evidence regarding the intervention is limited.

Table 5.8. Quality of evidence

	Quality of evidence
<i>Benefits of intervention:</i>	
Baseline health data	Medium-Strong
Relative risks (RR) for health benefits	Medium-Strong
Valuation of mortality	Medium-Strong
Productivity benefits (time savings)	Limited
<i>Cost of intervention:</i>	
Cost of ending OD (foregone benefit)	Limited
Total evidence	Limited

Source: Author.

6. Conclusion

This paper has evaluated the benefits and costs of four household drinking water and sanitation interventions. The interventions were selected on the basis of coverage rates and behavioral dimensions in the state. The benefit-cost ratios (BCRs) are found to be the largest for improved urban drinking water supply and improved rural sanitation, followed by improved urban sanitation, household point-of-use (POU) treatment of drinking water and improved rural water supply.

The quality of evidence associated with these interventions range from “medium” to “medium-strong”. A behavioral change campaign (BCC) promoting the use of existing sanitation by all household members has a smaller BCR, and supported by only “limited” quality of evidence.

Table 6.1. Summary of the benefits and costs and interventions (Rs million annualized)

	Interventions	Discount	Benefit	Cost	BCR	Quality of Evidence
1a	Improved drinking water supply - urban	3%	21,144	1,968	10.7	Medium
		5%	20,294	2,280	8.9	
		8%	19,127	2,797	6.8	
1b	Improved drinking water supply - rural	3%	18,929	3,801	5.0	Medium
		5%	18,167	4,403	4.1	
		8%	17,120	5,402	3.2	
2	BCC promotion of household POU treatment of drinking water*	3%	23,195	3,347	6.9	Medium-Strong
		5%	23,136	3,422	6.8	
		8%	23,050	3,538	6.5	
3a	Improved sanitation - urban	3%	34,941	5,422	6.4	Medium-Strong
		5%	33,527	5,441	6.2	
		8%	31,584	5,518	5.7	
3b	Improved sanitation - rural	3%	215,880	25,998	8.3	Medium-Strong
		5%	207,151	26,683	7.8	
		8%	195,164	27,987	7.0	
4	BCC promotion of use of existing sanitation facilities*	3%	1,243	671	1.9	Limited
		5%	1,192	673	1.8	
		8%	1,123	679	1.7	

Notes: All figures use VSL for valuation of mortality benefits and cost-of-illness (COI) approach for valuation of morbidity benefits. * "Mid" scenario. Source: Author.

References

- Andres, L., Briceno, B., Chase, C., and Echenique, J. (2014). Sanitation and Externalities: Evidence from Early Childhood Health in Rural India. Policy Research Working Paper 6737. World Bank. Washington, DC. USA.
- Black, R., Allen, L., Bhutta, Z., et al. (2008). Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*, 371, pp. 243-60.
- Borghini, J., Guinness, L., Ouedraogo, J., and Curtis, V. (2002). Is Hygiene Promotion Cost-Effective? A Case Study in Burkino Faso. *Tropical Medicine and International Health*, 7(11).
- Fewtrell, L., A. Prüss-Üstün, R. Bos, F. Gore, and Bartram, J. (2007). *Water, Sanitation and Hygiene: Quantifying the Health Impact at National and Local Levels in Countries with Incomplete Water Supply and Sanitation Coverage*. Environmental Burden of Disease Series 15. Geneva: World Health Organization.
- Fishman, M.S., L.E. Caulfield, M. De Onis, M. Blossner, A.A. Hyder, L. Mullany, and Black, R.E. (2004). *Childhood and Maternal Underweight*. In Ezzati, M., A.D. Lopez, A. Rodgers, and C.J.L. Murray (Eds): *Comparative Quantification of Health Risks – Global and Regional Burden of Disease Attributable to Selected Major Risk Factors*. Vol. 1. Geneva: World Health Organization.
- IIPS. 2017. National Family Health Survey IV 2015-16. State Fact Sheet Andhra Pradesh. International Institute for Population Sciences. Mumbai, India.
- IIPS. 2008. National Family Health Survey III 2005-06 Andhra Pradesh. International Institute for Population Sciences. Mumbai, India.
- Larsen, B. (2007). *Cost of environmental health risk in children under five years of age: Accounting for malnutrition in Ghana and Pakistan*. Background report prepared for the World Bank. Environment Department, World Bank.
- MSP/GOI. 2017. Swachhta Status Report 2016. Ministry of Statistics and Programme Implementation. Government of India. New Delhi, India.
- Olofin I, McDonald CM, Ezzati M, et al. (2013). Associations of suboptimal growth with all-cause and cause-specific mortality in children under five years: a pooled analysis of ten prospective studies. *PLoS ONE* 2013; 8: e64636.
- Pinfold, J. and Horan, N. (1996). Measuring the Effect of a Hygiene Behaviour Intervention by Indicators of Behaviour and Diarrhoeal Disease. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, Vol 90.
- Pruss-Ustun, A., Bartram, J., Clasen, T., Colford, J., et al. (2014). Burden of disease from inadequate water, sanitation and hygiene in low- and middle-income settings: a retrospective analysis of data from 145 countries. *Trop Med Int Health*, 19, pp. 894-905.

Saade, C., Bateman, M., and Bendahmane, D. (2001). *The Story of A Successful Public-Private Partnership in Central America: Handwashing for Diarrheal Disease Prevention*. Published by BASICS II, EHP, Unicef, USAID, and World Bank.

UNICEF (2018). *Financial and Economic Impacts of the Swachh Bharat Mission in India*. The United Nations Children's Fund. New Delhi. India.

Waddington, H., Snilstveit, B., White, H., and Fewtrell, L. (2009). *Water, sanitation and hygiene interventions to combat childhood diarrhoea in developing countries*. Synthetic Review 001. International Initiative for Impact Evaluation (3ie).

Wolf, J., Pruss-Ustun, A., Cumming, O., Bartram, J., et al. (2014). Assessing the impact of drinking water and sanitation on diarrhoeal disease in low- and middle-income settings: systematic review and meta-regression. *Tropical Medicine and International Health*, 19, pp. 928-942.

World Bank (2008). *Environmental health and child survival: Epidemiology, economics, experiences*. Washington, DC: World Bank.

World Bank (2016). *Methodology for valuing the health impacts of air pollution: Discussion of challenges and proposed solutions*. World Bank. Washington DC. USA.

Annex 1. Health effects of unsafe drinking water and sanitation

Unsafe drinking water and sanitation (WAS) is directly and indirectly affecting population health. Directly, unsafe WAS causes diarrheal infections and other health effects which in turn lead to mortality especially in young children. Indirectly, unsafe WAS contributes to poor nutritional status in young children through the effect of repeated diarrheal infections (World Bank, 2008; Fewtrell et al, 2007; Larsen, 2007).¹³ Poor nutritional status in turn increases the risk of child mortality or fatality rate from disease (Fishman et al., 2004; Black et al, 2008; Olofin et al, 2013).

Direct effect

Pruss-Ustun et al (2014), based on a global review, presents a methodology for estimating the diarrheal disease burden from unsafe WAS and the disease burden that can be expected to be prevented from improved WAS. For drinking water, the (preventable) disease burden is estimated relative to unimproved drinking water sources. For POU treatment (filtering or boiling), the (preventable) burden is estimated in relation to type of drinking water supply. For sanitation, a distinction is only made between unimproved and improved sanitation facilities as studies of health risks of other sanitation dimensions are insufficient. The relative risks of disease that are relevant for this paper are presented in tables A1.1-2.

For improved sanitation, however, a study of rural households in India by Andres et al (2014) estimates a relative risk of 0.53 that includes both the direct health benefit of improved sanitation to the household as well as the external health benefit to a community with high sanitation coverage rates.

Table A1.1. Relative risk of diarrheal disease and mortality for drinking water source and POU treatment

Baseline drinking water source	Improved drinking water source other than piped to premise	Basic piped water to premise	POU treatment (filtering or boiling and safe storage)
Unimproved drinking water source	0.89	0.77	0.55
Improved drinking water source other than piped to premise		0.86	0.62
Basic piped water to premise			0.72

Source: Pruss-Ustun et al (2014).

Table A1.2. Relative risk of diarrheal disease and mortality for sanitation

Baseline sanitation	Improved sanitation facility
Unimproved sanitation facility	0.72

Source: Pruss-Ustun et al (2014); Wolf et al (2014).

Indirect effect

¹³ Repeated infections, and especially diarrheal infections, have been found to significantly impair weight gains in young children. Studies documenting and quantifying this effect have been conducted in communities with a wide range of infection loads in a diverse group of countries. World Bank (2008) provides a review of these studies.

Estimating the indirect mortality effects of diarrhea from WAS is here undertaken in two stages. First, the fraction of under-five child mortality attributable to child underweight is estimated. This follows the methodology in Olofin et al (2013). Second, a fraction of under-five child mortality from underweight is attributed to diarrheal infections from WAS in early childhood using the approach in Fewtrell et al (2007).¹⁴

Estimates of increased risk of cause-specific mortality in children under five years of age with mild, moderate and severe underweight is presented in table A1.3 based on Olofin et al (2013).

Table A1.3. Relative risk of mortality from severe, moderate and mild underweight in children under five

	Severe	Moderate	Mild	None
Diarrhea	11.56	2.86	1.73	1.00
Acute lower respiratory infections (ALRI)	10.10	3.11	1.85	1.00
Measles	7.73	3.12	1.00	1.00
Malaria	1.29	1.65	1.26	1.00

Source: Olofin et al (2013). ALRI is acute lower respiratory infections. Relative risks are in relation to severe, moderate and mild underweight according to the WHO Child Growth Standards.

These relative risk ratios are applied to prevalence rates of child underweight to estimate attributable fractions (AF_j) of mortality by cause, j , from child underweight as follows:

$$AF_j = \frac{\sum_{i=1}^n P_i (RR_{ji} - 1)}{\sum_{i=1}^n P_i (RR_{ji} - 1) + 1} \quad (A1.1)$$

where RR_{ji} is relative risk of mortality from cause “ j ” for children in each of the underweight categories, i , in table A1.3; and P_i is the underweight prevalence rate.

Annual cases of mortality from child underweight by cause, “ j ” are estimated as follows:

$$M_j = BDB_j * AF_j \quad (A1.2)$$

where BDB_j is baseline annual cases of mortality from cause “ j ”. Annual cases of under-five child mortality from diarrheal infections in early childhood (W) is then estimated as follows:

$$W = \sum_{j=1}^{j=m} \gamma_j M_j \quad (A1.3)$$

¹⁴ An alternative approach to estimating the fraction of mortality attributable to diarrheal infections from WAS is the methodology developed in Larsen (2007) and World Bank (2008). This, however, requires estimation of counterfactual prevalence rates of child underweight (prevalence of underweight in the absence of diarrheal infections) from original survey data of child nutritional status. As the original survey data are often not readily available, the approach in Fewtrell et al is here used instead. The approach in Fewtrell et al gives a somewhat lower estimate of indirect mortality from WAS than the Larsen and World Bank methodology.

where γ_j is the fraction of child underweight mortality (M_j) attributed to diarrheal infections in early childhood. A value $\gamma_j = 0.5$ for ALRI, measles, and malaria is applied here based on Fewtrell et al (2007).

Preventable disease burden from interventions

The potential impact fraction (PIF) formula is applied to estimate the preventable disease burden from the drinking water and sanitation interventions evaluated in this paper:

$$PIF = \frac{\sum_{i=1}^n P_i RR_i - \sum_{i=1}^n P'_i RR_i}{\sum_{i=1}^n P_i RR_i} \quad (A1.4)$$

where P_i is the pre-intervention population distribution by type of drinking water, POU treatment, or sanitation; P'_i is the post-intervention population distribution; and RR_i is the relative risk of disease and mortality in tables A1.1-2.

The preventable disease burden or health benefit (B) of interventions is then:

$$B = PIF * BDB \quad (A1.5)$$

where B is annual cases of mortality and morbidity prevented, BDB is the baseline disease burden or annual cases of diarrheal disease mortality and morbidity in Andhra Pradesh. Included in the BDB in this paper is also annual cases of typhoid and paratyphoid mortality and annual cases of mortality (W) from the indirect effect discussed above.

As a new state, Andhra Pradesh faces a bright future, but it is still experiencing many acute social and economic development challenges. It has made great strides in creating a positive environment for business, and was recently ranked 2nd in India for ease of doing business. Yet, progress needs to be much faster if it is to achieve its ambitions of becoming the leading state in India in terms of social development and economic growth. With limited resources and time, it is crucial that focus is informed by what will do the most good for each rupee spent. The Andhra Pradesh Priorities project as part of the larger India Consensus – a partnership between Tata Trusts and the Copenhagen Consensus Center, will work with stakeholders across the state to identify, analyze, rank and disseminate the best solutions for the state. We will engage people and institutions from all parts of society, through newspapers, radio and TV, along with NGOs, decision makers, sector experts and businesses to propose the most relevant solutions to these challenges. We will commission some of the best economists in India, Andhra Pradesh, and the world to calculate the social, environmental and economic costs and benefits of these proposals



ANDHRA PRADESH PRIORITIES

AN
INDIA CONSENSUS
PRIORITIZATION
PROJECT

For more information visit www.APpriorities.com

C O P E N H A G E N C O N S E N S U S C E N T E R

Copenhagen Consensus Center is a think tank that investigates and publishes the best policies and investment opportunities based on social good (measured in dollars, but also incorporating e.g. welfare, health and environmental protection) for every dollar spent. The Copenhagen Consensus was conceived to address a fundamental, but overlooked topic in international development: In a world with limited budgets and attention spans, we need to find effective ways to do the most good for the most people. The Copenhagen Consensus works with 300+ of the world's top economists including 7 Nobel Laureates to prioritize solutions to the world's biggest problems, on the basis of data and cost-benefit analysis.